PROTEIN **POST-TRANSLATIONAL** MODIFICATION

INTRODUCTION OF POST-TRANSLATIONAL MODIFICATIONS

Living systems are inundated with diverse intracellular and environmental signals. Therefore, cells must be able to rapidly respond to programmed and unprogrammed cues.

Post-translational modification (PTM) refers to the covalent, usually enzymatic modification of proteins, and protein process during or after protein biosynthesis. PTM provides an energy-saving mechanism that can reversibly regulate protein functions in a rapid time frame. Protein PTM increases the functional diversity of the proteome by modifying proteins with functional groups such as phosphate, acetate, amide or methyl, and affects almost all aspects of normal cell biology and pathogenesis. It plays a key role in many cellular processes, such as cell differentiation, protein degradation, signal transduction and regulation processes, gene expression regulation, and protein-protein interactions.

Common types of protein post-translational modifications

- ① Disulfde Bonds
 - Linkage between two Cys residues, mostly intramolecular
 - Stabilizes protein structure
 - Currently no large-scale enrichment method available

1

6

Ν

- ⁽²⁾ Ubiquitination (Ubq)
 - Ubiquitin, a 8.5 kDa protein, is attached to Lys via an isopeptide bond
 - Mediates protein degradation, signaling, trafficking
 - Upon tryptic digestion, a Gly-Gly residue remains at the Lys and modified peptides can be enriched using anti-diGly-antibodies

Ubiquitin

④ Methylation

Ubiquitin

2 Ubiquitin 1 - Attachment of a methyl С group to Lys/Arg Κ - Regulates RNA processing, CH₃ 3 transcription, DNA damage ⁶ N-/O- Glycosylation 4 repair, protein translocation - Attachment of glycans to Acetyl R/K (mostly Arg), and epigenetic Asn or Ser/Thr/Tyr residues ③ Acetylation regulation of gene transcrip-- Structure as information - Attachment of an acetyl S,T, ticn carrier group to Lys - Enrichment using antibodies - Predominantly on extra-5 - Mediates protein-protein cellular domains involved interaction in cell-cell interaction and - Enrichment using acetylrecoghition, involved in ⁽⁵⁾ Phosphorylation lysine-specfic antibodies

protein folding and chaperone recruitment - Participates in many disease processes and is a biomarker for the diagnosis of multiple diseases - Multiple enrichment methods

- Combined structure and site-specific glycoproteomics remains challenging

X≠P S,T

- Attachment of a phosphate group to Ser/Thr/Tyr hydroxyl residues

- Enzyme activation switch, activity regulation

- Multiple enrichment methods

Acetyl

7

Signal Peptide

- ⑦ N-Term-Acetylation
 - Acetylation on mature protein N-termini, often after removal of initial Met
 - Linked to protein stability, localization and interaction
 - Enrichment using SCX, COFRADIC, TAILS, ChaFRADIC

STRATEGIES TO ANALYZE PROTEIN PTM





REFERENCES

Pagel O, Loroch S, Sickmann A, et al. Current strategies and findings in clinically relevant post-translational modification-specific proteomics. Expert review of proteomics, 2015, 12(3): 235-253.

Krishna R G, Wold F. Post-translational modifications of proteins//Methods in protein sequence analysis. Springer, Boston, MA, 1993: 167-172.

Prus G, Hoegl A, Weinert B T, et al. Analysis and interpretation of protein post-translational modification site stoichiometry. Trends in biochemical sciences, 2019, 44(11): 943-960.





© Creative Proteomics All Rights Reserved.